

CULPEPER SURGERY CENTER

Colon Cancer... You Can Prevent It

America's #2 Cancer Killer

- Colorectal cancer is the number 2 cancer killer in the United States, yet it is one of the most preventable types of cancer. Colorectal cancer is often curable when detected early.

Risk Factors

- Lifetime risk of colorectal cancer is roughly equal in men and women.
- Colorectal cancer is most common after age 50, but it can strike at younger ages. The risk of developing colorectal cancer increases with age.



Symptoms

Most early colorectal cancers produce no symptoms. This is why screening for colorectal cancer is so important. Some possible symptoms, listed below, do not always indicate the presence of colorectal cancer, but should prompt a visit with your physician and a check-up:

- New onset of abdominal pain
- Blood in or on the stool
- A change in stool caliber or shape
- A change in typical bowel habits, constipation, diarrhea

Who is Considered High Risk?

Colonoscopy is recommended for individuals of any age who are at higher than average risk for developing colorectal cancer by virtue of:

- Personal history of colorectal cancer or colorectal polyps
- A strong family history of the disease
- Inherited forms of colorectal polyps or cancer
- Predisposing chronic digestive condition such as inflammatory bowel disease (Crohn's disease or ulcerative colitis)

Recommendations for how often colonoscopy should be performed vary for different subsets of high risk individuals, and they should consult with their physician.



Colonoscopy: Preferred Screening Strategy


Colonoscopy is the preferred method of screening for colorectal cancer. The American College of Gastroenterology considers colonoscopy the "gold standard" for colorectal screening because colonoscopy allows physicians to look directly at the entire colon and to identify suspicious growths. Colonoscopy is the only test that allows a biopsy or removal of a polyp at the very same time it is first identified.



Colorectal Screening for African Americans

African Americans are diagnosed with colorectal cancer at a younger age than other ethnic groups, and African Americans with colorectal cancer have decreased survival compared with other ethnic groups.

Physician experts from the American College of Gastroenterology in 2005 issued new recommendations to healthcare providers to begin colorectal cancer screening in African Americans at age 45 rather than 50 years.



American College of Gastroenterology
Digestive Disease Specialists Committed to Quality in Patient Care

www.acg.gi.org

Please talk to your Primary Care Provider about your risks and need for a screening colonoscopy. The Specialists at the Culpeper Surgery Center will work with your Providers to take care of you.

541 Sunset Lane, Culpeper VA 22701
phone 540-829-0700 • fax 540-829-8191
www.culpepersurgerycenter.com

Providing outstanding ambulatory surgical care for the community

